

**Reading assignment**

Paul, Whites, and Nasar, 2.8, 2.9, 2.12 -&gt; 2.14

**Software**

div\_curl\_example.m

**Problem 1 - Line integrals & curl**


The magnetic field of a straight wire of radius  $a$  which has a constant current density  $J_0$ , is given by:

$$\mathbf{B} = \mu_0 J_0 r / 2 \mathbf{a} \quad \text{inside the wire } (r < a)$$

$$\mathbf{B} = \mu_0 J_0 a^2 / (2 r) \mathbf{a} \quad \text{outside the wire } (r > a).$$

where  $\mu_0$  and  $J_0$  are constants.

- a. Calculate  $\oint \mathbf{B} \cdot d\mathbf{l}$  around the 2 paths shown in the figure below. (The drawing shows a view as if the wire had been cut).



- b. Calculate  $\nabla \times \mathbf{B}$  for both regions.

**Problem 2 - Properties of fields with curl**

The electric field created by a cylinder of radius  $a$  with constant charge density  $\rho_0$  is:

$$\mathbf{E} = \rho_0 r / (2 \epsilon_0) \mathbf{a}_r \quad \text{inside the cylinder } (r < a) \text{ and}$$

$$\mathbf{E} = \rho_0 a^2 / (2 \epsilon_0 r) \mathbf{a}_r \quad \text{outside the cylinder } (r > a).$$

where  $\rho_0$  and  $\epsilon_0$  are constants.

- a. Verify that  $\oint \mathbf{E} \cdot d\mathbf{l} = 0$  on the same path as above and that  $\nabla \times \mathbf{E} = 0$ .  
 b. An illustration of the  $\mathbf{E}$  and  $\mathbf{B}$  fields can be obtained by running div\_curl\_example.m using matlab. Fig. 1 is the  $\mathbf{B}$  field while Figure 3 is the  $\mathbf{E}$  field. What are the properties of a field with non-zero curl?

**Problem 3 - Stokes theorem**

Calculate  $\oint (\nabla \times \mathbf{B}) \cdot d\mathbf{s}$  over the surface area enclosed by the 2 paths in Problem 1 (the shaded area). Compare your answer with the result from Problem 1a.

## Gradient, Line integrals, &amp; Curl

**Problem 4 - Gradient**

Compute the gradient of the following functions.

a.  $f = 8a^2 \cos \theta + 2rz$  (cylindrical)

b.  $f = a \cos^2 \theta / r$  (spherical)

Use the worksheet associated with Problem 2.8.1 in "Visual Electromagnetics for Mathcad" to check your answer. (You may have to use a specific number instead of the variable  $a$ ).

c. Calculate  $\nabla \times f$  for each of the functions above.